

# Calculus @ QFinance

## Lesson 11

Tuesday October 9<sup>th</sup> 2012

Fubini theorem

professor Daniele Ritelli

[www.unibo.it/faculty/daniele.ritelli](http://www.unibo.it/faculty/daniele.ritelli)



## Product of $\sigma$ -algebras

Let  $(\Omega_1, \mathcal{A}_1, \mu_1)$ ,  $(\Omega_2, \mathcal{A}_2, \mu_2)$  two measure spaces.

Put  $\Omega := \Omega_1 \times \Omega_2$ .

We want to build a measure  $\mu$  on  $\Omega$  which agrees with the given measures on  $\Omega_1$  and  $\Omega_2$ . Thus we have to provide the domain, i.e. a suitable  $\sigma$ -algebra on  $\Omega$ .

The product  $\sigma$ -algebra  $\mathcal{A}$  turns out to be the minimum  $\sigma$ -algebra on  $\Omega$  which contains rectangles  $A_1 \times A_2$  with  $A_1 \in \mathcal{A}_1$  and  $A_2 \in \mathcal{A}_2$ . So this  $\sigma$ -algebra is generated by the collection of rectangular sets

$$\mathcal{R} = \{A_1 \times A_2 \mid A_1 \in \mathcal{A}_1, A_2 \in \mathcal{A}_2\}$$

## Theorem

Product  $\sigma$ -algebra  $\mathcal{A}_1 \times \mathcal{A}_2$  is the minimum  $\sigma$ -algebra such that projections

$$\text{Pr}_1 : \Omega \rightarrow \Omega_1, \quad \text{Pr}_1(\omega_1, \omega_2) = \omega_1$$

$$\text{Pr}_2 : \Omega \rightarrow \Omega_2, \quad \text{Pr}_2(\omega_1, \omega_2) = \omega_2$$

are measurable

The most relevant situation for application is when  $\Omega_1 = \Omega_2 = \mathbb{R}$ ,  $\mathcal{A}_1 = \mathcal{A}_2 = \mathcal{B}$  Borel  $\sigma$ -algebra in  $\mathbb{R}$ .

Borel set in the plane can be generated in two equivalent ways.

The most relevant situation for application is when  $\Omega_1 = \Omega_2 = \mathbb{R}$ ,  $\mathcal{A}_1 = \mathcal{A}_2 = \mathcal{B}$  Borel  $\sigma$ -algebra in  $\mathbb{R}$ .

Borel set in the plane can be generated in two equivalent ways.

### Theorem

$\sigma$ -algebras generated by

$$\mathcal{R} = \{B_1 \times B_2 \mid B_1, B_2 \in \mathcal{B}\}, \quad \mathcal{I} = \{I_1 \times I_2 \mid I_1, I_2 \text{ intervals}\}$$

are the same.

**Product measure: Guido Fubini Venice 1879, New York 1943**

To build product measure we need to work with  $\sigma$ -finite measure space  $(\Omega_1, \mathcal{A}_1, \mu_1)$ ,  $(\Omega_2, \mathcal{A}_2, \mu_2)$ .

## Product measure: Guido Fubini Venice 1879, New York 1943

To build product measure we need to work with  $\sigma$ -finite measure space  $(\Omega_1, \mathcal{A}_1, \mu_1)$ ,  $(\Omega_2, \mathcal{A}_2, \mu_2)$ . This is the case for Lebesgue measure and for probability measures. We will use  $\mu$  to indicate product measure on product  $\sigma$ -algebra  $\mathcal{A}_1 \times \mathcal{A}_2$ .

## Product measure: Guido Fubini Venice 1879, New York 1943

To build product measure we need to work with  $\sigma$ -finite measure space  $(\Omega_1, \mathcal{A}_1, \mu_1)$ ,  $(\Omega_2, \mathcal{A}_2, \mu_2)$ . This is the case for Lebesgue measure and for probability measures. We will use  $\mu$  to indicate product measure on product  $\sigma$ -algebra  $\mathcal{A}_1 \times \mathcal{A}_2$ .

First step: we impose “natural” condition:

$$\mu(A_1 \times A_2) = \mu(A_1)\mu(A_2)$$



## Product measure: Guido Fubini Venice 1879, New York 1943

To build product measure we need to work with  $\sigma$ -finite measure space  $(\Omega_1, \mathcal{A}_1, \mu_1)$ ,  $(\Omega_2, \mathcal{A}_2, \mu_2)$ . This is the case for Lebesgue measure and for probability measures. We will use  $\mu$  to indicate product measure on product  $\sigma$ -algebra  $\mathcal{A}_1 \times \mathcal{A}_2$ .

First step: we impose “natural” condition:

$$\mu(A_1 \times A_2) = \mu(A_1)\mu(A_2)$$

To assign a measure to non rectangular sets we need to introduce the notion of **section** of a subset  $A$  of  $\Omega_1 \times \Omega_2$

If  $A \subset \Omega_1 \times \Omega_2$  and if  $\omega_2 \in \Omega_2$  section of foot  $\omega_2$  is the subset of  $\Omega_1$

$$A_{\omega_2} = \{\omega_1 \in \Omega_1 \mid (\omega_1, \omega_2) \in A\}$$

If  $A \subset \Omega_1 \times \Omega_2$  and if  $\omega_2 \in \Omega_2$  section of foot  $\omega_2$  is the subset of  $\Omega_1$

$$A_{\omega_2} = \{\omega_1 \in \Omega_1 \mid (\omega_1, \omega_2) \in A\}$$

If  $A \subset \Omega_1 \times \Omega_2$  and if  $\omega_1 \in \Omega_1$  section of foot  $\omega_1$  is the subset of  $\Omega_2$

$$A_{\omega_1} = \{\omega_2 \in \Omega_2 \mid (\omega_1, \omega_2) \in A\}$$

If  $A \subset \Omega_1 \times \Omega_2$  and if  $\omega_2 \in \Omega_2$  section of foot  $\omega_2$  is the subset of  $\Omega_1$

$$A_{\omega_2} = \{\omega_1 \in \Omega_1 \mid (\omega_1, \omega_2) \in A\}$$

If  $A \subset \Omega_1 \times \Omega_2$  and if  $\omega_1 \in \Omega_1$  section of foot  $\omega_1$  is the subset of  $\Omega_2$

$$A_{\omega_1} = \{\omega_2 \in \Omega_2 \mid (\omega_1, \omega_2) \in A\}$$

## Theorem

If  $A \in \mathcal{A}_1 \times \mathcal{A}_2$  then for any  $\omega_2 \in \Omega_2$  we have that  $A_{\omega_2} \in \mathcal{A}_1$  and for any  $\omega_1 \in \Omega_1$  we have that  $A_{\omega_1} \in \mathcal{A}_2$

## Definition

If  $A \in \mathcal{A}_1 \times \mathcal{A}_2$  define

$$\mu(A) = \int_{\Omega_2} \mu_1(A_{\omega_2}) \, d\mu_2(\omega_2) \quad (\star)$$

## Definition

If  $A \in \mathcal{A}_1 \times \mathcal{A}_2$  define

$$\mu(A) = \int_{\Omega_2} \mu_1(A_{\omega_2}) \, d\mu_2(\omega_2) \quad (\star)$$

## Theorem

If  $\mu_1, \mu_2$  are  $\sigma$ -finite measures, then functions

$$\omega_2 \mapsto \mu_1(A_{\omega_2}), \quad \omega_1 \mapsto \mu_2(A_{\omega_1})$$

are, respectively measurable with respect to  $\mathcal{A}_2, \mathcal{A}_1$  and

## Definition

If  $A \in \mathcal{A}_1 \times \mathcal{A}_2$  define

$$\mu(A) = \int_{\Omega_2} \mu_1(A_{\omega_2}) \, d\mu_2(\omega_2) \quad (\star)$$

## Theorem

If  $\mu_1, \mu_2$  are  $\sigma$ -finite measures, then functions

$$\omega_2 \mapsto \mu_1(A_{\omega_2}), \quad \omega_1 \mapsto \mu_2(A_{\omega_1})$$

are, respectively measurable with respect to  $\mathcal{A}_2, \mathcal{A}_1$  and

$$\int_{\Omega_1} \mu_2(A_{\omega_1}) \, d\mu_1(\omega_1) = \int_{\Omega_2} \mu_1(A_{\omega_2}) \, d\mu_2(\omega_2)$$

## Theorem

Set function  $\mu$  introduced in  $(\star)$  is a measure. It is **unique** since any other measure which coincides with  $\mu$  on rectangles is equal to  $\mu$  on product  $\sigma$ -algebra  $\mathcal{A}_1 \times \mathcal{A}_2$



## Fubini Theorem on nested integrals

If  $f \in \mathcal{L}^1(\Omega_1 \times \Omega_2)$  then functions

$$\omega_1 \mapsto \int_{\Omega_2} f(\omega_1, \omega_2) d\mu_2(\omega_2), \quad \omega_2 \mapsto \int_{\Omega_1} f(\omega_1, \omega_2) d\mu_1(\omega_1)$$

belong respectively to  $\mathcal{L}^1(\Omega_1)$ ,  $\mathcal{L}^1(\Omega_2)$  and the following formula holds

## Fubini Theorem on nested integrals

If  $f \in \mathcal{L}^1(\Omega_1 \times \Omega_2)$  then functions

$$\omega_1 \mapsto \int_{\Omega_2} f(\omega_1, \omega_2) \, d\mu_2(\omega_2), \quad \omega_2 \mapsto \int_{\Omega_1} f(\omega_1, \omega_2) \, d\mu_1(\omega_1)$$

belong respectively to  $\mathcal{L}^1(\Omega_1)$ ,  $\mathcal{L}^1(\Omega_2)$  and the following formula holds

$$\int_{\Omega_1 \times \Omega_2} f(\omega_1, \omega_2) \, d(\mu_1 \times \mu_2)(\omega_1, \omega_2) =$$

## Fubini Theorem on nested integrals

If  $f \in \mathcal{L}^1(\Omega_1 \times \Omega_2)$  then functions

$$\omega_1 \mapsto \int_{\Omega_2} f(\omega_1, \omega_2) d\mu_2(\omega_2), \quad \omega_2 \mapsto \int_{\Omega_1} f(\omega_1, \omega_2) d\mu_1(\omega_1)$$

belong respectively to  $\mathcal{L}^1(\Omega_1)$ ,  $\mathcal{L}^1(\Omega_2)$  and the following formula holds

$$\begin{aligned} \int_{\Omega_1 \times \Omega_2} f(\omega_1, \omega_2) d(\mu_1 \times \mu_2)(\omega_1, \omega_2) = \\ \int_{\Omega_1} \left( \int_{\Omega_2} f(\omega_1, \omega_2) d\mu_2(\omega_2) \right) d\mu_1(\omega_1) = \end{aligned}$$

## Fubini Theorem on nested integrals

If  $f \in \mathcal{L}^1(\Omega_1 \times \Omega_2)$  then functions

$$\omega_1 \mapsto \int_{\Omega_2} f(\omega_1, \omega_2) \, d\mu_2(\omega_2), \quad \omega_2 \mapsto \int_{\Omega_1} f(\omega_1, \omega_2) \, d\mu_1(\omega_1)$$

belong respectively to  $\mathcal{L}^1(\Omega_1)$ ,  $\mathcal{L}^1(\Omega_2)$  and the following formula holds

$$\begin{aligned} \int_{\Omega_1 \times \Omega_2} f(\omega_1, \omega_2) \, d(\mu_1 \times \mu_2)(\omega_1, \omega_2) &= \\ \int_{\Omega_1} \left( \int_{\Omega_2} f(\omega_1, \omega_2) \, d\mu_2(\omega_2) \right) d\mu_1(\omega_1) &= \\ \int_{\Omega_2} \left( \int_{\Omega_1} f(\omega_1, \omega_2) \, d\mu_1(\omega_1) \right) d\mu_2(\omega_2) \end{aligned}$$

## Integration in Euclidean spaces. Notations

$\mathbb{R}^m = \mathbb{R}^p \times \mathbb{R}^q$  with  $p + q = m$

$(x, y) \in \mathbb{R}^m$  means  $x \in \mathbb{R}^p$  e  $y \in \mathbb{R}^q$ .

## Integration in Euclidean spaces. Notations

$\mathbb{R}^m = \mathbb{R}^p \times \mathbb{R}^q$  with  $p + q = m$

$(x, y) \in \mathbb{R}^m$  means  $x \in \mathbb{R}^p$  e  $y \in \mathbb{R}^q$ .

If  $A \subset \mathbb{R}^m$  and  $x \in \mathbb{R}^p$ , the  **$x$  section of  $A$**  is the subset  $A_x$  in  $\mathbb{R}^q$  defined by:

$$A_x := \{y \in \mathbb{R}^q \mid (x, y) \in A\}$$

## Integration in Euclidean spaces. Notations

$\mathbb{R}^m = \mathbb{R}^p \times \mathbb{R}^q$  with  $p + q = m$


$(x, y) \in \mathbb{R}^m$  means  $x \in \mathbb{R}^p$  e  $y \in \mathbb{R}^q$ .

If  $A \subset \mathbb{R}^m$  and  $x \in \mathbb{R}^p$ , the  **$x$  section of  $A$**  is the subset  $A_x$  in  $\mathbb{R}^q$  defined by:

$$A_x := \{y \in \mathbb{R}^q \mid (x, y) \in A\}$$

If  $y \in \mathbb{R}^q$  the  **$y$  section of  $A$**  is defined by:

$$A_y := \{x \in \mathbb{R}^p \mid (x, y) \in A\}$$





## Section's Theorem

Let  $A \subset \mathbb{R}^m$  measurable. Then if  $x \in \mathbb{R}^p$ , section  $A_x$  is a.e. measurable. Moreover  $x \mapsto \ell_q(A_x)$  is a measurable function and

$$\ell_m(A) = \int_{\mathbb{R}^p} \ell_q(A_x) d\ell_p(x)$$

$\ell_m$  stands for Lebesgue measure in  $\mathbb{R}^m$ . The same for  $\ell_p$ ,  $\ell_q$

**Fubini's Theorem** G. Fubini: “Il teorema di riduzione per gli integrali doppi” Rend. Sem. Mat. Torino, vol.9, 1949.

**Fubini's Theorem** G. Fubini: “Il teorema di riduzione per gli integrali doppi” Rend. Sem. Mat. Torino, vol.9, 1949.

Assume  $A \subset \mathbb{R}^m$  measurable and let  $f \in \mathcal{L}(A)$ . Define  $\mathcal{S}$  as the subset of  $\mathbb{R}^p$  where all  $q$ -section of  $A$  have positive measure

$$\mathcal{S} = \{x \in \mathbb{R}^p \mid \ell_q(A_x) > 0\}$$


**Fubini's Theorem** G. Fubini: “Il teorema di riduzione per gli integrali doppi” Rend. Sem. Mat. Torino, vol.9, 1949.

Assume  $A \subset \mathbb{R}^m$  measurable and let  $f \in \mathcal{L}(A)$ . Define  $\mathcal{S}$  as the subset of  $\mathbb{R}^p$  where all  $q$ -section of  $A$  have positive measure

$$\mathcal{S} = \{x \in \mathbb{R}^p \mid \ell_q(A_x) > 0\}$$


Then the following equality holds

$$\int_A f(x, y) dx dy = \int_{\mathcal{S}} \left( \int_{A_x} f(x, y) dy \right) dx$$



**Example.** If  $A = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq x \leq 1, x^2 \leq y \leq x + 1\}$  evaluate:

$$\iint_A xy \, dx \, dy$$



$$\iint_A xy \, dx \, dy = \int_0^1 \left( \int_{x^2}^{1+x} xy \, dy \right) dx$$



$$\iint_A xy \, dx \, dy = \int_0^1 \left( \int_{x^2}^{1+x} xy \, dy \right) dx = \int_0^1 x \left[ \frac{y^2}{2} \right]_{x^2}^{1+x} dx$$

$$\iint_A xy \, dx \, dy = \int_0^1 \left( \int_{x^2}^{1+x} xy \, dy \right) dx = \int_0^1 x \left[ \frac{y^2}{2} \right]_{x^2}^{1+x} dx$$
$$\iint_A xy \, dx \, dy = \frac{1}{2} \int_0^1 (-x^5 + x^3 + 2x^2 + x) \, dx$$


$$\iint_A xy \, dx \, dy = \int_0^1 \left( \int_{x^2}^{1+x} xy \, dy \right) dx = \int_0^1 x \left[ \frac{y^2}{2} \right]_{x^2}^{1+x} dx$$
$$\iint_A xy \, dx \, dy = \frac{1}{2} \int_0^1 (-x^5 + x^3 + 2x^2 + x) \, dx = \frac{5}{8}$$

## Example

Given  $A = \{(x, y) \in \mathbb{R}^2 \mid y \geq 0, y \leq -x + 3, y \leq 2x + 3\}$ . Evaluate:

$$\iint_A y \, dx \, dy$$

Integration domain: triangle with vertices in  $(-\frac{3}{2}, 0)$ ,  $(3, 0)$ ,  $(0, 3)$ .



We first integrate in  $x$  and then in  $y$  :

$$\int_0^3 \left( \int_{\frac{y-3}{2}}^{3-y} y \, dx \right) dy$$

We first integrate in  $x$  and then in  $y$  :

$$\int_0^3 \left( \int_{\frac{y-3}{2}}^{3-y} y \, dx \right) dy = \int_0^3 \left( \frac{9}{2}y - \frac{3}{2}y^2 \right) dy$$

We first integrate in  $x$  and then in  $y$  :


$$\int_0^3 \left( \int_{\frac{y-3}{2}}^{3-y} y \, dx \right) dy = \int_0^3 \left( \frac{9}{2}y - \frac{3}{2}y^2 \right) dy = \frac{27}{4}$$


## Example

Evaluate the measure of

$$A = \left\{ (x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1 \right\} \cap \left\{ (x, y) \in \mathbb{R}^2 \mid 0 \leq y \leq x \leq \sqrt{3} y \right\}$$







$$\int_0^{1/\sqrt{2}} \left( \int_{x/\sqrt{3}}^x dy \right) dx + \int_{1/\sqrt{2}}^1 \left( \int_{x/\sqrt{3}}^{\sqrt{1-x^2}} dy \right) dx$$

For exercise. The only difficult point is

$$\int \sqrt{1 - x^2} \, dx = \frac{1}{2} \left( x \sqrt{1 - x^2} + \arcsin x \right)$$

Two integers  $a$  and  $b$  are said to be coprime if the only positive integer that evenly divides both of them is 1. This is the same thing as their greatest common divisor  $\text{gcd}(a, b) = 1$ .

Two integers  $a$  and  $b$  are said to be coprime if the only positive integer that evenly divides both of them is 1. This is the same thing as their greatest common divisor  $\text{gcd}(a, b) = 1$ .

**Theorem** (E. Cesàro 1881) The probability that two randomly chosen numbers are coprime is given by a product over all primes:

$$\prod_{p \in \mathbb{P}} \left( 1 - \frac{1}{p^2} \right) = \frac{1}{\sum_{n=1}^{\infty} \frac{1}{n^2}}$$

Two integers  $a$  and  $b$  are said to be coprime if the only positive integer that evenly divides both of them is 1. This is the same thing as their greatest common divisor  $\text{gcd}(a, b) = 1$ .

**Theorem** (E. Cesàro 1881) The probability that two randomly chosen numbers are coprime is given by a product over all primes:

$$\prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^2}\right) = \frac{1}{\sum_{n=1}^{\infty} \frac{1}{n^2}}$$

G.H. Hardy; E. M. Wright (2008). An Introduction to the Theory of Numbers (6th ed.) Oxford University Press, theorem 332.

**Theorem** (L. Euler)

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

PROOF. (D. Ritelli: to appear on American Mathematical Monthly  
mid 2013)

**Theorem** (L. Euler)

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

PROOF. (D. Ritelli: to appear on American Mathematical Monthly mid 2013)

Let  $A = [0, \infty) \times [0, \infty)$  consider

$$\iint_A \frac{dx dy}{(1+y)(1+x^2 y)}$$



First integrate with respect to  $x$  and then with respect to  $y$  finding

$$\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} = \int_0^\infty \left( \frac{1}{1+y} \int_0^\infty \frac{dx}{1+x^2 y} \right) dy$$

First integrate with respect to  $x$  and then with respect to  $y$  finding

$$\begin{aligned}\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} &= \int_0^\infty \left( \frac{1}{1+y} \int_0^\infty \frac{dx}{1+x^2 y} \right) dy \\ &= \int_0^\infty \left( \frac{1}{1+y} \left[ \frac{\arctan(\sqrt{y} x)}{\sqrt{y}} \right]_{x=0}^{x=\infty} \right) dy\end{aligned}$$

First integrate with respect to  $x$  and then with respect to  $y$  finding

$$\begin{aligned}\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} &= \int_0^\infty \left( \frac{1}{1+y} \int_0^\infty \frac{dx}{1+x^2 y} \right) dy \\ &= \int_0^\infty \left( \frac{1}{1+y} \left[ \frac{\arctan(\sqrt{y} x)}{\sqrt{y}} \right]_{x=0}^{x=\infty} \right) dy \\ &= \frac{\pi}{2} \int_0^\infty \frac{dy}{\sqrt{y}(1+y)} = \frac{\pi}{2} \int_0^\infty \frac{2u}{u(1+u^2)} du = \frac{\pi^2}{2}\end{aligned}$$

Then we proceed reverting the order of integration:

$$\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} = \int_0^\infty \left( \int_0^\infty \frac{dy}{(1+y)(1+x^2 y)} \right) dx$$

Then we proceed reverting the order of integration:

$$\begin{aligned}\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} &= \int_0^\infty \left( \int_0^\infty \frac{dy}{(1+y)(1+x^2 y)} \right) dx \\ &= \int_0^\infty \frac{1}{1-x^2} \left( \int_0^\infty \left( \frac{1}{1+y} - \frac{x^2}{1+x^2 y} \right) dy \right) dx\end{aligned}$$

Then we proceed reverting the order of integration:

$$\begin{aligned}\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} &= \int_0^\infty \left( \int_0^\infty \frac{dy}{(1+y)(1+x^2 y)} \right) dx \\ &= \int_0^\infty \frac{1}{1-x^2} \left( \int_0^\infty \left( \frac{1}{1+y} - \frac{x^2}{1+x^2 y} \right) dy \right) dx \\ &= \int_0^\infty \frac{1}{1-x^2} \left[ \ln \frac{1+y}{1+x^2 y} \right]_{y=0}^{y=\infty} dx\end{aligned}$$

Then we proceed reverting the order of integration:

$$\begin{aligned}\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} &= \int_0^\infty \left( \int_0^\infty \frac{dy}{(1+y)(1+x^2 y)} \right) dx \\&= \int_0^\infty \frac{1}{1-x^2} \left( \int_0^\infty \left( \frac{1}{1+y} - \frac{x^2}{1+x^2 y} \right) dy \right) dx \\&= \int_0^\infty \frac{1}{1-x^2} \left[ \ln \frac{1+y}{1+x^2 y} \right]_{y=0}^{y=\infty} dx \\&= \int_0^\infty \frac{1}{1-x^2} \ln \frac{1}{x^2} dx\end{aligned}$$

Then we proceed reverting the order of integration:

$$\begin{aligned}\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} &= \int_0^\infty \left( \int_0^\infty \frac{dy}{(1+y)(1+x^2 y)} \right) dx \\&= \int_0^\infty \frac{1}{1-x^2} \left( \int_0^\infty \left( \frac{1}{1+y} - \frac{x^2}{1+x^2 y} \right) dy \right) dx \\&= \int_0^\infty \frac{1}{1-x^2} \left[ \ln \frac{1+y}{1+x^2 y} \right]_{y=0}^{y=\infty} dx \\&= \int_0^\infty \frac{1}{1-x^2} \ln \frac{1}{x^2} dx \\&= \int_0^\infty \frac{\ln x^2}{x^2 - 1} dx =\end{aligned}$$



Then we proceed reverting the order of integration:

$$\begin{aligned}\iint_A \frac{dx dy}{(1+y)(1+x^2 y)} &= \int_0^\infty \left( \int_0^\infty \frac{dy}{(1+y)(1+x^2 y)} \right) dx \\&= \int_0^\infty \frac{1}{1-x^2} \left( \int_0^\infty \left( \frac{1}{1+y} - \frac{x^2}{1+x^2 y} \right) dy \right) dx \\&= \int_0^\infty \frac{1}{1-x^2} \left[ \ln \frac{1+y}{1+x^2 y} \right]_{y=0}^{y=\infty} dx \\&= \int_0^\infty \frac{1}{1-x^2} \ln \frac{1}{x^2} dx \\&= \int_0^\infty \frac{\ln x^2}{x^2 - 1} dx = \\&= 2 \int_0^\infty \frac{\ln x}{x^2 - 1} dx\end{aligned}$$

Hence, equating we get

$$\int_0^\infty \frac{\ln x}{x^2 - 1} dx = \frac{\pi^2}{4}. \quad (\text{a})$$

Hence, equating we get

$$\int_0^{\infty} \frac{\ln x}{x^2 - 1} dx = \frac{\pi^2}{4}. \quad (\text{a})$$

Now split the integration domain in (a) between  $[0, 1]$  and  $[1, \infty)$  and change the variable  $x = 1/u$  in the second integral, so that

$$\begin{aligned} \int_0^{\infty} \frac{\ln x}{x^2 - 1} dx &= \int_0^1 \frac{\ln x}{x^2 - 1} dx + \int_1^{\infty} \frac{\ln x}{x^2 - 1} dx \\ &= \int_0^1 \frac{\ln x}{x^2 - 1} dx + \int_0^1 \frac{\ln u}{u^2 - 1} du. \end{aligned} \quad (\text{b})$$

From (a) and (b) we get

$$\int_0^1 \frac{\ln x}{x^2 - 1} dx = \frac{\pi^2}{8}. \quad (c)$$

From (a) and (b) we get

$$\int_0^1 \frac{\ln x}{x^2 - 1} dx = \frac{\pi^2}{8}. \quad (c)$$

But we have already shown that

$$\int_0^1 \frac{\ln x}{x^2 - 1} dx = \sum_{n=0}^{+\infty} \frac{1}{(2n+1)^2} \quad (d)$$

From (a) and (b) we get

$$\int_0^1 \frac{\ln x}{x^2 - 1} dx = \frac{\pi^2}{8}. \quad (c)$$

But we have already shown that

$$\int_0^1 \frac{\ln x}{x^2 - 1} dx = \sum_{n=0}^{+\infty} \frac{1}{(2n+1)^2} \quad (d)$$

Hence from (c) and (d) follows

$$\sum_{n=0}^{+\infty} \frac{1}{(2n+1)^2} = \frac{\pi^2}{8}$$

To get Euler's statement define  $E = \sum_{n=1}^{+\infty} \frac{1}{n^2}$  and split the series in considering even and odd indexes:

$$\sum_{n=1}^{\infty} \frac{1}{(2n)^2} + \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = E$$

To get Euler's statement define  $E = \sum_{n=1}^{+\infty} \frac{1}{n^2}$  and split the series in considering even and odd indexes:

$$\sum_{n=1}^{\infty} \frac{1}{(2n)^2} + \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = E$$

Observe that  $\sum_{n=1}^{\infty} \frac{1}{(2n)^2} = \sum_{n=1}^{\infty} \frac{1}{4n^2} = \frac{E}{4}$



To get Euler's statement define  $E = \sum_{n=1}^{+\infty} \frac{1}{n^2}$  and split the series in considering even and odd indexes:

$$\sum_{n=1}^{\infty} \frac{1}{(2n)^2} + \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = E$$

Observe that  $\sum_{n=1}^{\infty} \frac{1}{(2n)^2} = \sum_{n=1}^{\infty} \frac{1}{4n^2} = \frac{E}{4}$  thus  $\sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = \frac{3}{4}E$

To get Euler's statement define  $E = \sum_{n=1}^{+\infty} \frac{1}{n^2}$  and split the series in considering even and odd indexes:

$$\sum_{n=1}^{\infty} \frac{1}{(2n)^2} + \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = E$$

Observe that  $\sum_{n=1}^{\infty} \frac{1}{(2n)^2} = \sum_{n=1}^{\infty} \frac{1}{4n^2} = \frac{E}{4}$  thus  $\sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = \frac{3}{4}E$  hence

$$\frac{3}{4} \sum_{n=1}^{+\infty} \frac{1}{n^2} = \frac{\pi^2}{8} \implies \sum_{n=1}^{+\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

Euler 1735: Basel Problem